


ARO 15529.12-GS

AD-A134664

Reprinted from MONTHLY WEATHER REVIEW, Vol. 111, No. 5, May 1983
American Meteorological Society
Printed in U. S. A.

**Duration of Convective Events Related to Visible Cloud, Convergence, Radar and Rain
Gage Parameters in South Florida**

RONALD L. HOLLE AND ANDREW I. WATSON



DTIC FILE COPY

83 11 08 066

Duration of Convective Events Related to Visible Cloud, Convergence, Radar and Rain Gage Parameters in South Florida¹

RONALD L. HOLLE AND ANDREW I. WATSON

Weather Research Program, Environmental Research Laboratories, NOAA, Boulder, CO 80303

(Manuscript received 2 September 1981, in final form 20 January 1983)

ABSTRACT

The time interval between initiation of surface convergence and the subsequent response of visible cloud growth to this convergence was examined for nine cases of convection that occurred over the FACE 1973 and 1975 mesonetworks in south Florida. Clouds ranged in size from small echoes with a few towers to merged lines or large clusters of towers, but they met a series of observational criteria that specified them as belonging to a similar set of clouds, and were not representative of the entire range of clouds in the area. Visible clouds first formed 10 to 55 min after the associated surface convergence began, and grew rapidly upward 20 to 100 min after convergence started.

This highly variable response could be understood better by taking into account the duration of the cloud, which is defined as the time from first surface convergence to complete dissipation. The same nine cases were examined as were chosen initially for the visible cloud study. When duration was considered, first visible cloud response occurred at an average of 15% through the cloud duration, and rapid upward cloud growth at 36%. Other parameters derived from divergence, radar- and gage-measured rainfall also tended to cluster within specific portions of the total duration of the cloud. The data for each event for the nine clouds are presented and described in terms of the cloud duration.

1. Introduction and background

In the past, photogrammetric studies have related visible cloud growth to radar, gage and synoptic data. However, there appears to be no earlier examination of the time interval between the start of surface convergence and the development of new visible clouds forming directly over the same convergence region. Plank (1969) related Florida cloud population distributions to peninsular-scale patterns, but not on a cloud-by-cloud basis. Orville (1965) studied the initiation of individual cumuli with stereo photogrammetry and attributed their growth to forcing by mountain ridges interacting with the ambient flow, but had no surface convergence data. Other researchers have used photogrammetry to consider various aspects of cloud growth, but not in relation to surface convergence, due to a lack of photo or wind data on the cloud scale. A study of the role of surface convergence in forcing clouds directly overhead has been performed with data collected during the Florida Area Cumulus Experiment (FACE) and is part of this report.

The original purpose of the research was to investigate the linkage between convergence and cloud for-

mation during the early stages of cloud growth. However, in the initial phase of the study, a large variation was found in this time interval, by up to a factor of five. The research was then expanded, in the second phase of the study, to seek a reason for the variation.

It was found that the highly variable time interval could be explained by considering the duration of the same cumulus systems as the clouds that were chosen for their visible cloud views. Short convergence/cloud-response times were associated with short-lived cloud systems, and long intervals related to long-lived systems. The shorter-lived cloud systems were smaller and less intense than the long-lasting ones. This second portion of the research goes beyond the Thunderstorm Project (Braham, 1952), where cumulative rainfall amount was shown to be associated with cumulative storm duration in a regular fashion during a storm's lifetime. From the FACE mesonetwork, a larger variety of data specific to the cloud itself is examined by relating several cloud development milestones to duration of the cloud systems.

2. Data and procedures

The data used in the study were obtained during the Florida Area Cumulus Experiment (FACE) in the southern portion of the peninsula. All cases but one were from FACE 1975; the data were collected in the mesonetwork shown in Fig. 1. Time-lapse cameras in 1975 were located at the three Doppler sites and

¹ A portion of this research was conducted while the authors were at the National Hurricane and Experimental Meteorology Laboratory, Coral Gables, FL 33146.




FIG. 1. FACE 1975 mesonetwork. Time-lapse cameras were located at all three Doppler radar sites and at the Field Observing Site (FOS). The wind network covered 1440 km².

the Field Observing Site (FOS) in the west-central network. One case was from FACE 1973 when the mesonetwork was located somewhat east of, but overlapping, the 1975 area. Hand-held photos but no time-lapse pictures were taken for the 1973 case. The Miami WSR-57 radar is located about 100 km from the center of the mesonetwork. The rain gages and wind stations in 1975 are shown in Fig. 1.

Time-lapse photographs in 1975 were taken on 16-mm color film looking toward the center of the mesonetwork at a rate of one frame per 5–7 seconds. For the study of visible cloud response to convergence, the first step was to identify periods when a significant but isolated new cloud formed and grew in the cameras' views. Next, the radar data were searched for an isolated echo that was strong enough to be identified with a distinct convergence center and visible cloud. In most situations, the cameras were not directed toward the exact location where the isolated echo was starting to grow, other clouds obscured the view, the cameras were not operating, or, most commonly, the very early stage of cloud formation was not identifiable or capable of being related simply to a surface convergence center. For large clouds, the cloud field was often too distant from the camera to

be certain of the location, and therefore a correspondence with the radar or convergence event could not be made at the early stage of new growth which was needed to define the start of surface convergence. For small clouds, the area of visible clouds during the early growth stage was not large enough to be detected readily with the wind station spacing (6.4 km average).

There were nine cases, however, where subject clouds grew in the mesonetwork and could be included in the sample. Smaller clouds in this set had several definable towers and single-cored radar echoes throughout their lifetime. Larger clouds were merged lines or clusters of towers which grew into echoes with several cores. In all cases, there was a distinguishable convergence center, smaller than the 1440 km² mesonetwork area, which was separate from other convergence patterns in south Florida. If the convective entity was not separable with the radar or convergence data, the case was not included in this study. Convergence areas and radar echoes in these nine cases ranged in size from a small portion (~10%) to most of the total area of the 1440 km² mesonetwork. The nine echoes selected for this paper belong to the upper third of the size distribution of echoes in this

area; however, they do not include echoes comprising the largest 5% of the distribution (Wiggert *et al.*, 1981). Lopez (1978) has shown that the largest 5% of the echoes account for 80% of the rainfall in the eastern Atlantic. Since rainfall in both the eastern Atlantic and Florida is largely generated by convection, a similar distribution of rainfall amount versus echo size may also occur in the FACE network.

3. Visible cloud response to convergence

The growth response of visible clouds directly over the area where first convergence occurred was described in terms of three parameters called Events A, B, and C. Event A was the time of first convergence, Event B was the time of first visible cloud response, and Event C was the time of rapid upward growth of the visible cloud.

It is recognized that prior to Event A used in this study, other parameters could be measured that produce the first convergence which was considered the start of these systems. Pressure perturbations, outflows from other clouds, etc., could be detected prior to first convergence in some cases. Here, the research began with convergence for all cases regardless of its source.

First convergence (Event A) was determined to the nearest 5 min from several of the products available for studies of convergence and rainfall on these days. First, the daily time profiles of area average divergence and weighted convergence (defined in Watson *et al.*, 1981) for the full 1440 km² mesonetwork were examined to find the 5-min period when the background, prevailing convergence increased to a stronger convergence value which was identifiable as related to a specific convective entity. Watson *et al.* showed that an increase in convergence of $25 \times 10^{-6} \text{ s}^{-1}$ in 10 min over the total network is the crucial parameter to relate to rainfall, and that concept was used for the larger and longer-lasting clouds whose convergence covered a significant portion of the network. Second, for the smaller clouds, area divergence quantities and their changes were calculated at 5-min intervals in the specific portion of the network where the cloud was located, such as a quadrant, because total network data were not sufficiently sensitive to isolate the cloud. Third, maps of divergence and streamlines over the mesonetwork were made at 5-min intervals to specify the time and locate the regions of development of small clouds. The combination of these 3 steps to define the start of a convergence event was not difficult to use for determining Event A for the nine cases. All other cases were omitted when the start of convergence was unclear from the data.

Visible cloud initiation (Event B) was often rather easy to determine to the nearest 5 min from the time-lapse film. In several cases, there were no clouds in

the surface convergence area; the time of first cumulus appearance was considered to be the response to the prior convergence. In the other situations, a disorganized and essentially random field of small non-raining, shallow cumulus clouds grew horizontally and vertically to become organized and clustered in a rather short time; cloud bases merged and became noticeably darker and harder at this stage. Some of these changes were from random cumuli to a new line, and others were to a new cluster of cloud elements. The area of the visible cloud initially was smaller, in general, than the convergence area over which the cloud formed. Later, as the cloud became mature or was dissipating, the mature cloud's radar echo grew to an area similar to the area of the originating convergence zone, as shown in the case study of Holle and Maier (1980).

The time of rapid upward visible cloud growth (Event C) was also rather easily found from the same film. It can best be described as the time when several or many towers are simultaneously growing upward very rapidly. At this time the cloud line or complex is passing into the stage where it is apparent that the cloud definitely will produce rainfall. Event C was found to the nearest 5 min.

Table 1 shows results of the analysis of nine clouds whose initial growth stages were well defined. For estimated (Est.) times, the discrepancy usually was ± 5 min from the given time. Data for the FACE 1973 case of June 15 were compiled from the analysis by Holle and Maier; the FACE 1975 case of August 19 has been the subject of research by Cuning *et al.* (1982). The interval from first convergence (Event A) to the time of initial organization of the visible cloud system (Event B) ranged from 10 to 55 min and av-

TABLE 1. Times of Events A, B, and C for nine cloud entities in FACE 1973 and 1975 mesonetworks. Events B and C are expressed in min after Event A.

Date	Time of first convergence	Visible cloud response	Rapid upward growth of visible clouds
	Event A	Event B (min)	Event C (min)
15 June 1973	1425 EDT	Est. 35	Est. 50
08 Aug. 1975	1645	Est. 20	45
12 Aug. 1975	Est. 1720	20	60
13 Aug. 1975	Est. 1425	Est. 25	75
18 Aug. 1975	Est. 1345	15	70
19 Aug. 1975	1450	55	100
20 Aug. 1975	Est. 1545	25	75
25 Aug. 1975 (A)	1305	Est. 20	45
25 Aug. 1975 (B)	Est. 1415	10	20
Average time (min)	0	25	60
Normalized time (%)	0	15	36

eraged 25 min, with a standard deviation of 11 min. It was another 35 min until the rapid upward growth stage (Event C) was reached, which was 60 min on the average after Event A (standard deviation of 13 min). The interval between convergence and visible cloud growth would be shorter for clouds that never produced radar echoes. During the growth stage of the nine subject clouds, variability in the time intervals can be partially explained by such factors as the rate of increase in convergence while the cloud grows (Watson *et al.*, 1981). They also found that convergence changes of a given amount produced different cloud responses when stratifications were made by humidity aloft, stability, and low-level wind speed.

4. Time history of storm events normalized to duration

The preceding lags of visible cloud growth after the initiation of surface convergence varied from case to case by as much as a factor of 5. Not shown in Table 1 is the fact that longer time intervals tended to be associated with longer cloud lifetimes and larger storms. To take this effect into account, the times of Events B and C were normalized to the total duration of the storm system. Event A was chosen as the start of the system's lifetime (0%), and the time of dissipation (100%), called Event J, was found from radar or time-lapse photos. When the nine cases were normalized on this scale, Event B occurred at an average of 15% (bottom line, Table 1) through the lifetime of the cloud entity, and Event C was at 36%.

It was apparent from these results for visible clouds that normalizing the events to cloud duration was useful in understanding the variations in time lags between the individual cases. Other specific events (Table 2) were then sought which could be identified for the same nine cases that were studied for visible cloud response. Since surface wind, rain gage and radar data were readily available over the duration of cloud entities originally identified for the visible cloud lag study, six more parameters were chosen to relate to duration for the second part of the research. These milestones were shown to occur in a similar order for south Florida convection in the conceptual model proposed by Ulanski and Garstang (1978) over a range of cell sizes. Here the investigation will be on the relationship between the time of these events relative to the duration of the system; no direct predecessor of this study is apparent. Two wind-related milestones were chosen: times of maximum convergence (Event E) and maximum divergence (Event I) associated with the cloud system. They were determined with the same types of information as that given earlier for first convergence (Event A). Another pair of events was derived from the Miami radar: times of first radar returns from the subject cloud (Event D) and maximum radar-estimated rainfall

TABLE 2. Events used to study storm duration in FACE mesonetworks.

Event	Description	Data
A	First convergence above background levels	Mesonetwork winds
B	Visible clouds first appear or are no longer randomly distributed	Time lapse photos
C	Visible clouds start rapid upward growth	Time lapse photos
D	First radar returns from cloud entity	Radar
E	Maximum convergence at cloud entity	Mesonetwork winds
F	First rain on ground	Rain gages and/or photos
G	Maximum radar rainfall from cloud entity	Radar
H	Maximum gage-measured rain from cloud	Rain gages
I	Maximum divergence at cloud	Mesonetwork winds
J	Complete dissipation	Radar and/or photos

from the cloud entity (Event G). The latter was a volume measure for the entire echo; the time of the maximum was for the entire echo found from 1) the time history graph of radar reflectivity for the entire mesonetwork, which may include other echoes, and 2) from maps showing the magnitude of the relevant echo's core intensity at 5-min intervals. The center of the radar beam was typically about 1 km above the surface in the mesonetwork. Finally, rain gage data were considered for two situations. Event F refers to the time of the first rain on the ground, which was detected either by gages or from time-lapse photos of the cloud. Subject clouds occurred principally over the uniform gage network (Fig. 1) so that gage density was not critical for finding Event F. Event H was the time when gages measured the maximum rainfall from the cloud (15-min accuracy at times), although there were several instances when clouds moved out of the gage and camera network. Finally, Event J is the time of complete dissipation from radar or photos.

Normalized times for Events A to J are listed in Table 3 and diagrammed in Fig. 2. Average duration (Events A-J) for the nine cloud entities was 161 min. The individual Events B and C, referring to visible cloud development, are shown by Table 1 in minutes and by Table 3 in normalized times. The first radar returns (Event D) also occurred 36% through the cloud system, the same as Event C, and in some situations occurred earlier than Event C. (Numbers in

TABLE 3. Normalized times, as percent of total duration, when events A to J occurred for nine cloud entities in FACE. Times are normalized to storm duration (right column), which extended from Event A as 0% to Event J as 100%. Normalized averages and standard deviations are at bottom of table.

Date	Event										Duration (min)
	A	B	C	D	E	F	G	H	I	J	
15 June 73	0	29	38	54	(38)	58	58	69	77	100	130
8 August 75	0	14	31	(24)	34	38	45	41-52	59	100	145
12 August 75	0	12	38	(36)	(34)	41	62	missing	69	100	160
13 August 75	0	14	42	(41)	(67)	50	72	69-78	86	100	180
18 August 75	0	9	44	(34)	47	(41)	47	56-66	75	100	160
19 August 75	0	23	43	45	51	51	(62)	55-57	81	100	235
20 August 75	0	13	39	39	39	41	62	(54-62)	69	100	195
25 August 75 (A)	0	15	33	33	(22)	41	81	missing	(56)	100	135
25 August 75 (B)	0	9	18	(14)	27	32	55	missing	59	100	110
Average (%)	0	15	36	36	40	44	60	61	70	100	161
σ (%) ($n-1$ method)	0	7	8	12	14	8	11	10	11	0	38

parentheses in Table 3 refer to events that were out of time order from the average sequence.) After another 4% of the duration (6 min), maximum convergence occurred (Event E), followed 6 min later by first rain on the ground (Event F); the times for events C, D, E and F are quite similar and are not deemed to be significantly different. Note that the time interval between first radar and first gage detection of rainfall averaged 8% of the duration, or 13 min. This time interval is similar to the results of Holle and Maier (1980), and extends the results of Watson *et al.* (1981) from their study of convergence using radar-estimated rainfall to results using gages. The mature stage

of the cloud systems is shown by the next three events (diagrammed as descending in Fig. 2). On the average, maximum radar-estimated rainfall (Event G) was at about the same time (60%) as maximum gage-measured rainfall (Event H) at 61%. Maximum surface divergence (Event I) occurred at 70% of the storm's duration. The dissipation stage from that point to the complete disappearance of the cloud entity (Event J) was another 30% of the cloud lifetime, compared with 15% from first convergence to first presence or organization of visible clouds. In summary, the highly variable times found from the study of visible clouds, and shown in Table 3 for the other events, are found




FIG. 2. Normalized cloud duration in % for Events A to J during nine cases of cloud entities growing in FACE mesonetwork during June 1973 and August 1975. Mean % of each event at center of diamond; standard deviation shaded to either side.

to be rather well organized by taking duration into account. The times when the events occur are not distributed uniformly over the total duration of the cloud, but tend to cluster in a specific portion of the normalized cloud duration, and this tendency helps explain the large variability in the time intervals before normalization.

5. Discussion

There were two major conclusions from this study. First, visible cloud development lagged convergence initiation over a wide range of time intervals. The average was 25 min from first convergence to first organization of the cloud field, and another 35 min passed, on the average, until the clouds began a rapid upward growth stage. This is apparently the first study to quantify these time intervals. Second, these variable time responses could be explained partially by taking into account the duration of the entire cloud system. Short periods from convergence initiation to an event such as visible cloud formation, rainfall or radar echo initiation tended to be associated with short times until the demise of the cloud system, and conversely for long periods and long-lived systems.

The lag of visible cloud development after convergence was sought from an additional source of time-lapse photography in a mesonet network in Illinois during the 1979 VIN program (University of Virginia-Illinois State Water Survey-NOAA). It proved impossible to define these early events sufficiently well to add to the sample presented here for FACE. The principal difficulties were advection into the mesonet of existing clouds, convergence apparently aloft rather than at the surface, large and complex convergence and cloud fields, and motion of growing clouds across the camera view. The events that were identified usually occurred in the same order in Illinois as in Florida; however, there were no cloud sequences that could be followed completely in the VIN network to comprise additional cases for the present study.

This study of visible cloud reaction to surface convergence has led to a more general examination of cloud milestones related to duration. Specific events within a cloud lifetime tended to cluster within rather narrow portions of the total duration of a cloud. How long these clouds lasted also was well-correlated

($r = .92$) with the rainfall production by the cloud entities, although this sample consists of a narrow range of sizes and types of clouds, as detailed earlier. Based on this study, the lag of visible clouds behind surface convergence has been specified for a small number of south Florida clouds, and should be studied elsewhere to see how well the conclusions apply. In addition, the role of duration in delineating the timing of cloud development should be examined further with larger data sets to find how well such features as the early stages of radar echoes or satellite images can provide information on how long the convective system will last.

Acknowledgments. Our thanks are extended to two NOAA employees, Mr. Tony Barnston and Mr. James Hansen, for advice and calculations related to this paper. This research was supported by the Atmospheric Research Section, National Science Foundation and the Army Research Office, Department of Defense, under NSF Grant ATM-78-08865.

REFERENCES

- Braham, R. R., 1952: The water and energy budgets of the thunderstorm and their relation to thunderstorm development. *J. Meteor.*, **9**, 227-242.
- Cunning, J. B., R. L. Holle, P. T. Gannon and A. I. Watson, 1982: Convective evolution and merger in the FACE experimental area: mesoscale convection and boundary layer interactions. *J. Appl. Meteor.*, **21**, 953-977.
- Holle, R. L., and M. W. Maier, 1980: Tornado formation from downdraft interaction in the FACE mesonet. *Mon. Wea. Rev.*, **108**, 1010-1028.
- Lopez, R. E., 1978: Internal structure and development processes of C-scale aggregates of cumulus clouds. *Mon. Wea. Rev.*, **106**, 1488-1494.
- Orville, H. D., 1965: A photogrammetric study of the initiation of cumulus clouds over mountainous terrain. *J. Atmos. Sci.*, **22**, 700-709.
- Plank, V. G., 1969: The size distribution of cumulus clouds in representative Florida populations. *J. Appl. Meteor.*, **8**, 46-67.
- Ullanski, S., and M. Garstang, 1978: The role of surface divergence and vorticity in the life cycle of convective rainfall. Part II: Descriptive model. *J. Atmos. Sci.*, **35**, 1063-1069.
- Watson, A. I., R. L. Holle, J. B. Cunning, P. T. Gannon and D. O. Blanchard, 1981: Low-level convergence and the prediction of convective precipitation in south Florida. Tech. Rep. No. 4, NOAA Environmental Research Laboratories, Boulder, CO, and Illinois State Water Survey, Champaign-Urbana, IL, 228 pp.
- Wiggert, V., G. J. Lockett and S. S. Ostlund, 1981: Radar rain-shower growth histories and variations with wind speed, echo motion, location and merger status. *Mon. Wea. Rev.*, **109**, 1467-1494.



DTIC COPY AVAILABLE	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947
---------------------------	---

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 15529.12-GS	2. GOVT ACCESSION NO. AN/A 134664	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) Duration of Convective Events Related to Visible Cloud, Convergence, Radar and Rain Gage Parameters in South Florida		5. TYPE OF REPORT & PERIOD COVERED Reprint
		6. PERFORMING ORG. REPORT NUMBER N/A
7. AUTHOR(s) Ronald L. Holle Andrew I. Watson		8. CONTRACT OR GRANT NUMBER(s) ARO 20-78
9. PERFORMING ORGANIZATION NAME AND ADDRESS Environmental Research Laboratories/NOAA Boulder, CO 80303		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS N/A
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE May 83
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Submitted for announcement only.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)